CIHM Microfiche Series (Monographs) ICMH
Collection de
microfiches
(monographies)



Canadian Institute for Historical Microreproductions / Institut canadian de microreproductions historiques

(C) 1997

Technical and Bibliographic Notes / Notes techniques et bibliographiques

copy may the slgni	Institute has attempt available for filming be bibliographically images in the reficantly change the ked below.	 Features of unique, which production, 	of this copy n may alter or which	which any of may	été plai ogra ou	possitire qui aphiqui qui pe normai	oie de sont p e, qui p uvent e e de fili	se proc eut-être peuvent exiger u mage so	urer. unlq mod ne m ont inc	Les dues du ifler un odlficat diqués	kemplali létalls de le point de le image tion dan ci-desso	e cet le vu repr	exem- e bibii- oduite,
Ш	Coloured covers / Couverture de coul	eur			<u>_</u>	1		ages / F					
	Covers damaged /					Pag	es dan	naged /	Pages	s endor	mmagée	S	
Ш	Couverture endom	nagée						ored and					
	Covers restored an	d/or laminated	d /			. ray	03 1034	aulees (evou	Pellicui	56 5		
	Couverture restaure	ée et/ou pellic	ulée		1			oloured olorées,			oxed / u plquée	s	
	Cover title missing	Le titre de co	ouverture ma	anque		, ,							
	Coloured maps / Ca	artes géograp	hiques en c	ouleur	L	Pag	es deta	iched / f	Pages	détaci	hées		
						Sho	wthroug	gh / Trai	nspar	ence			
	Coloured Ink (i.e. of												
	Encre de couleur (i.	e. autre que t	pieue ou noi	re)				rint vari		!			
	Coloured plates and	d/or illustration	ne /			Qua	iite ine	gale de	i'impr	ession			
	Planches et/ou illus					Inclu	ides su	ppleme du maté	ntary	materia	al /		
\Box	Bound with other m	aterial /					picha	ou maic	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	аррісііі	enane		
Ι Χ Ι	Relié avec d'autres	documents									red by o		
	Only edition availab Seule édition dispon					pos: parti	sible i elleme	mage nt obscu	/ Les	par un	es total feuillet d	leme l'errat	nt ou a, une
	Tight binding may ca interior margin / La							eilleure				GC 16	ayon a
	l'ombre ou de la d intérieure.	listorsion le l	ong de la r	marge		disc	olourati	ons are	filme	d twice	ng colo to ensu	re th	e best
	Blank leaves added within the text. Whe omitted from filming blanches ajoutée apparaissent dans l possible, ces pages	never possible / II se peut ques lors d'un e texte, mais,	e, these have ue certaines ne restaux lorsque cels	e been pages ration		colo	rations es deu	variab	les o	des (décolora la meille	ations	sont
	Additional commen												
	Commentaires supp	olémentaires:											
	tem is filmed at the redu cument est filmé au taux			ous.									
10x	14x		18x		22x			26x			30x		
		TIT			1					Т	T	Π	
	12v	16v		204			24 ×			200			

The copy filmed here has been reproduced thanks to the generosity of:

National Library of Canada

The images eppearing here are the best quality possible considering the condition end legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and anding on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shell contain the symbol —— (meening "CONTINUED"), or the symbol ∇ Imsening "END"), whichever applies.

Meps, platas, cherts, etc., may be filmed at different reduction ratios. Those too largs to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames se raquirad. The following diagrams illustrate the method:

L'axemplaire filme fut reproduit grâce à la génorosité de:

Bibliothèque nationale du Canada

Les images suiventes ont été reproduites avec le plus grand soin, compts tenu de la condition et de la netteté de l'exempleirs filmé, et en conformité evec les conditions du contret de filmege.

Les exempleires originaux dont le couverture en pepier est imprimée sont filmés en commençent par le premier plet et an terminent soit per le dernière paga qui comports una ampreinta d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres examplairas origineux sont filmés en commençent per la première page qui comporte une empreinta d'imprassion ou d'illustration et an terminant par la darnièra page qui comporte una telle empreinte.

Un des symboles suivents appereitra sur la dernièra imaga da chaqua microficha, selon ls cas: le symbole → signifis "A SUIVRE", la symbole ♥ signifia "FIN".

Lee certes, plenches, tableeux, atc., peuvent âtrs filmée à des taux de réduction différents.
Lorsque le document est trop grand pour être reproduit en un œut cliché, il eet filmé à pertir de l'engle supérieur gauche, de geuche à droits, et da haut an bes, en prenant le nombra d'imeges nécesseire. Les diagrammes suivants illustrent la méthode.


1 2 3	1	2	3
-------	---	---	---

1	
2	
3	

1	2	3
4	5	6

MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)






APPLIED IMAGE Inc

1653 East Main Street Rochester, New York 14609 USA (716) 482 - 0300 - Phone

(716) 288 - 5989 - Fax



DAIRY AND COLD STORAGE COMMISSIONER'S BRANCH OTTAWA, - - - CANADA


THE CARE OF CREAM FOR BUTTERMAKING

BY

GEO. H. BARR Chief, Dairy Division.

BULLETIN No. 32—DAIRY AND COLD STORAGE SERIES

Published by direction of the Hon. Martin Burrell, Minister of Agriculture, Ottawa, Ont.



LETTER OF TRANSMITTAL.

OTTAWA, February 20, 1912.

To the Honourable

The Minister of Agriculture.

SIR,—I have the honour to submit for your approval, the manuscript for a bulletin on 'The Care of Cream for Buttermaking,' which has been prepared under my direction by Mr. Geo. H. Barr, Chief of the Dairy Division of this Branch. The information contained herein is derived from the pages of a previous bulletin and from the results of recent investigations carried on by the Dairy Division. I beg to recommend that it be printed for distribution as Bulletin No. 32 of the Dairy and Cold Storage Series.

I have the honour to be, Sir.

Your obedient servant,

J. A. RUDDICK,

Dairy and Cold Storage Commissioner.

plan hav gett farn fact

Editorial and the second of th

the has qual as f

of t

and will one

THE CARE OF CREAM FOR BUTTERMAKING.

BY

Geo. H. Barr,

Chief of the Dairy Division.

The manufacture of butter in creameries on what is known as the gathered cream plan is becoming more and more general throughout Canada. The advantage of having fresh warm skimmilk for feeding purposes, the desire to avoid the risk of getting disease in their herds from the mixed skimmilk from a larger number of farms, together with a lower cost of hauling and, in most cases, a lower cost for manufacturing the butter, makes the system popular with many milk producers.

On the other hand, the fact must be recognized that in most cases a change from the system of separating the milk at the creamery to that of gathering the cream, has resulted in an inferior quality of butter being made, for the reason that the quality of the cream separated at the farms, when delivered at the creameries, is not as fine as that from milk separated at the creamery.

Cream which is separated on the farm can be delivered to the creamery in as mod condition as that which is separated at the creamery. It is simply a question of the patron who skims his milk at home taking proper care of the cream and having it delivered to the creamery often enough.

It is admitted by all creamery authorities that finer butter can be made from cream which is sweet when delivered at the creamery, than from cream which is sour and curdled. It is a well known that any taint that may be in the milk or cream will be to some expectations are carried into the butter. Therefore, the producer will see at since the responsibility resting upon him in securing a fine flavoured butter at the creamery.

In the production of fine flavoured cream, the same precautions must be observed as those which are necessary in furnishing milk to separator creameries or to cheese factories. The following are some of the essential points:—

Feeds that will Injure the Flavour of the Butter, and which should not be

Fed to Milch Cows.

- 1. Turnips and turnip lops.
- 2. Rape or rye.
- 3. Decayed ensilage.
- 4. Leeks, onions, or apples in large quantities.

Other Causes of Taints in Cream.

- 1. Cows' udders and teats in an unclean condition at milking time.
- 2. Milking in unclean stables.
- 3. Using unclean, wooden, galvanized or rusty milking pails.

- 4. Separating the milk in the stable.
- 5. Improperly cleaned separators.
- 6. Keeping the c. im in cellars or other places where there are roots or vegetables.
- 7. Keeping the cream for several days at a temperature over 55 degrees.
- 8. ('ows drinking water from stagnant ponds, or the leakage from barnyards.

Conditions that are Necessary to Produce Fine-flavoured Cream.

Pure Water.—The cows should have at all times an abundant supply of pure water to drink. When cows are compelled to drink the water of swamps, muddy ponds or sluggish streams and ditches, in which there is decaying animal matter, including their own droppings, there is a constant menace to their health, and unless the cows are in good health, they cannot give first-class milk. Moreover, the mud often full of foul germs, which collects on the legs, flanks and udders of the cows and falls into the milk at the time of milking, is a direct source of infection.

Salt.—When cows have free access to salt at all times, they will keep in better health, will give more milk, and the cream from this milk will have a better flavour, and keep sweet longer, than when they do not get any at all, or receive it only at intervals.

Milking.—Cleanliness in the stable is desirable at all times, but especially a milking time should the stables be clean and free from dust. The udders, teats and flanks of the cow should be brushed before milking. Only bright, clean tin pails should be used to milk in. Galvanized pails are difficult to keep clean, and bad flavours have been traced to their use.

The Hand Power Separator.

The hand power cream separator is the most reliable and best method of skimming milk at the farm, and the only method that can be recommended. Nearly all the separators on the market will do efficient skimming if properly handled.

Handling and Care of the Separator.—It is important that the separator runs smoothly. Any trembling or shaking of the separator while skimming will cause a loss of butter fat in the skimmilk. Only special separator oil should be used, and it is well to make a run about a ce in three weeks, using kerosene oil on all the bearings.

In skimming, three things must be observed: (1) The speed of the separator must be maintained according to the directions sent with it. The only reliable way to do this, is to count the number of revolutions of the crank by the watch. A low speed means loss of fat in the skimmilk. (2) The flow of the milk into the separator should be uniform. (3) The temperature of the milk should not be under 90 degrees, and for that reason, the best time to separate the milk is immediately after milking. A low temperature is also liable to cause loss of fat in the skimmilk. The faster the milk passes through the separator, the less complete is the separation, and a thinner cream is given. One of the questions often asked by patrens is: Why does my test vary so? When one knows that the speed of the machine, the flow of the





Fig. 1.



Fic 2



iles.

ure ddy ter, less rud,

ow: tter our.

r at ails ba-t

imall

un4 use and the

tot ray 017 20%

ng. ter a

the

milk, and the temperature of the milk all affect the test of the cream, it is not difficult to understand why it may vary considerably. A variation in the test does not necessarily mean any loss to the patron. Every separator has some device for changing the test of the cream. In most cases the adjustment is at the cream outlet. If so, by turning the cream serew in, the cream will be richer, and by turning it out, the cream will be thinner.

All the parts of the separator which come in contact with the milk or cream should be washed in luke warm water, to which has been added a small quantity of sal soda or other cleansing powder, and then thoroughly scalded with boiling water each time the separator is used.

In the

In inst

In inst

it had

would

more

in op

at the

cause the a

plain

vessel

handy

shoule

h cream

cream were

rancie T

Leng

1

cent a tempe

Location of Separator.—In some cases the separators are placed in the coverables. This may be a convenient arrangement, but it is not by any means a proper place for separating milk, unless a special room, well ventilated and lighted, is partitioned off to exclude the stable odours and dust. This room should have a smooth cement floor, which can be easily cleaned.

Advantages of a Rich Cream.—It is doubtful if there is any one thing which injures the quality of gathered cream butter so much as thin cream. Thin cream is responsible to a large extent for the old sour cream flavour so frequently found on gathered cream butter.

Many patrons have the idea that a large amount of cream should give a correspondingly large amount of money, forgetting that they are paid only for the butter fat in the cream, or the butter made from the fat.

Skimming a rich cream is a decided advantage to the patrons in more ways than one. The following table shows the advantage with reference to hauling and the amount of skimmilk retained on the farm, of skimming a rich eream with a herd of ten cows averaging 4,000 lbs. of 3.5 per cent milk.

TABLE I.-TOTAL MILK 40,000 LBS. TOTAL BUTTER FAT 1,400 LBS.

	Sent to the	Creamery.	Kept on the Farm.		
Per cent fat in cream.	Butter Fat.	Cream.	Skimmilk.	Skimmilk.	
20	Lbs. 1,400 1,400 1,400 1,400 1,400	Lbs. 7,000 5,600 4,662 4,004 3,500	Lbs. 33,000 34,400 35,338 35,996 36,500	Extra lbs. 1,400 2,338 2,996 3,500	

The foregoing table shows that there is the same amount of fat sent to the ereamery in all cases, but a very great difference in the amount of skimmilk left on the farm. An additional advantage is that there is less cream to take care of, and it is easier to keep it sweet.

Table II shows the results of experiments conducted by the Dairy Division, on keeping thick and thin cream from the same milk under exactly the same conditions for different periods.

TABLE II.

Cream kept.	Length of time kept.	Per cent fat in cream.	Temp. of cream.	Per cent acid in cream.
In the cellars		31·00 22·20	65·50 66·50	·430 ·540
n insulated tanks	36 hrs	31·67 22·00	57·50 57·50	· 175
in insulated tanks	60 hrs	32·19 21·55	55°40 55°50	*380 *440

These figures show that in every case the thin cream had the highest acidity, and it had always a much stronger flavour.

If all the cream sent to cream gathering creameries tested 30 per cent fat, it would mean thousands of dollars of extra money in the pockets of the patrons from more and better stock, and the quality of the butter would be very much improved.

Vessels for Holding Cream.—Many patrons keep the eream in earthen crocks, or in open pails. Crocks are liable to get broken or chipped, and experiments conducted at the O.A.C. Dairy School, Guclph, show that earthen crocks if chipped in any way cause an undesirable flavour in the butter. Cream kept in open pails is exposed to the air too much and, for that reason, is apt to become tainted. A well soldered plain bottomed tin can about 8 inches in diameter and 20 inches deep is the best-vessel in which to keep cream. This style of can (Fig. 1) is easy to keep clean and handy to put into a tank of water and ice. When two lots are mixed, the cream should be well stirred. Fig. 2 shows a very useful utensil for this purpose.

Where to Keep the Cream.

Keeping Cream in Cellars.—A great deal more than half of the cream sent to the creamerics is kept in cellars. Our own experiments proved that we could not keep cream sweet for thirty-six hours, or for delivery every other day, in cellars which were as cool as the ordinary run of farm house cellars; also that the cream when left uncovered developed a strong cellar flavour and the butter had a tendency to become rancid.

The following table shows the temperature, and acidity, by the acidimeter test, of cream kept in the cellars for different periods.

TABLE III.

Length of time kept.	Average temp. of cellars.	Average temp. of cream.	Averageper cent acid.
36 hrs.	63 7 deg.	64·5 deg.	:47
60 "	62 9 "	63·5 "	:50
72 "	64 0 "	64·0 "	:52

Note.—Freshly separated cream will have according to this test about 13 per cent acid; the acidity continues to increase more or less rapidly, according to the temperature. When it reaches 25 per cent, it begins to be perceptible to the taste,

and may be called 'sour.' The change is a gradual one, and the difference between what we call 'sweet' cream and 'sour' cream is one of degree only. The human sense of taste is not as delicate as the acidimeter test and cannot detect the same differences. When the cream reaches the stage of thickening, it has an acidity of about .35 per cent.

owe

that

no d

old.

:9.88

creat

> erei well wat

> hou

deg

hav

per

exp

Th

Keeping Cream in Water and Ice.—We have found that the easiest and best way to keep cream sweet and clean in flavour is to put it in a shot gun can (Fig. 1) and place it in a tank of water and ice immediately after skimming. We can recommend an insulated tank as shown in Fig. 3. This tank is made with a space of four inches filled with planing mill shavings on all sides and on the bottom, the cover also being insulated in the same manner. It is lined inside with galvanized iron. Such a tank is a little expensive, but it is certainly a great saver of ice. An ordinary wooden tank is next best. All tanks should have covers, as they help to keep down the temperature of the water and cream.

Table IV shows the advantage of keeping the cream in an insulated tank with water and ice, compared with the best results we could get by keeping it in the cellars. The cream was divided into two lots immediately after skimming, one lot set in the cellar and the other set in water and ice.

TABLE IV.

Treatment given the cream.	Length of time kept,	Average temp. of cream.	Average acidity of cream.
Cooled in tank	36 hrs. 36 " 60 "	52·8 64·5 53·0 63·7	157 1470 165 1505

These results show that the cellar cream had three times as much acid as that cooled in water and ice.

Keeping Cream in a Refrigerator.—Some people have recommended cooling the cream by placing it in a refrigerator immediately after skimming.

The following table shows the results of dividing cream equally into two lots cooling one lot in ice and water and the other in a first-class refrigerator.

TABLE V.

Cream kept in.	Average temperature	Average acidity.	Ice used,
Refrigerator		·415% ·230%	Lbs. 221 178

There were 43 lbs. more ice used in the refrigerator than in the water. The average temperature of the cream kept in the refrigerator was nearly one degree

lower, yet the acidity was almost twice as high. This is no doubt due to the fact that ice and water will cool the cream much faster than cold air.

en

an

me

27

nd nd

105

ng

th

The eovers were kept on the eream cans in both cases and there was practically difference in the flavour of the cream at any time.

Both lots of butter scored 42.5 points for flavour when fresh. When three weeks old, the butter from the water and ice cooled cream scored 40.77 points and the other \$3.88 points.

Table VI shows the effect that different temperatures have on the acidity of cream kept for different periods.

TABLE VI.

	Length	of time	kept.		Average t	emp. o	f cream.	Average per cent aci l
11 · · · · · · · · · · · · · · · · · ·	 				55 57 58 . 64 53	16 deg	•	145 170 190 210 510 150 310
н .	 		• • • • •	· · · · · · · · · · · · · · · · · · ·	47	5		· 165 · 390

It will be observed that when the eream was cooled to 55 degrees soon after skimming, it kept perfectly sweet for thirty-six hours, or for delivery to the creamery every other day. This temperature can be secured at most farms with the ordinary well water if an insulated tank is used. If this temperature cannot be secured with water alone, ice should be used.

To keep eream sweet for eighty-four hours, or for delivery twice a week, it must be kept down to 48 degrees. To do this, ice must be used. The lot kept eighty-four hours at 54 degrees was cooled in an insulated tank with water from the well at 48 degrees and changed night and morning. It was quite sour when delivered at the creamery

We were able to keep cream perfectly sweet for eighty-four hours, but it did not have the elean pleasant flavour which is found on cream kept sweet for shorter periods.

Keeping the cream for longer than two days at the farms has, no doubt, much to do with the old cream flavour so common in gathered eream butter, and we can searcely expect to have this defect remedied so long as cream is gathered less than three times each week.

When the eream is pasteurized at the creamery, the loss of butter fat in the buttermilk will be greater if the eream is sour than if delivered in a sweet condition. There is also a greater loss of fat in pasteurizing thin cream than thick. All cream separated at the farms should test between 27 and 35 per cent butter fat.

The Creamery Owner's Responsibility.

While asking the patrons to make improvement in their methods, we do not wish to relieve the ereamery owners and managers of their responsibilities to the patrons.

They must see to it that the equipment of the creamery is such that the cream supplied is handled in the most efficient manner; that the testing is done accurately and honestly, and that the creamery is a model of cleanliness and a standing object lesson for the patrons.

These conditions cannot be secured or maintained unless there is a reasonable price paid for manufacturing. Modern creamery equipment is expensive, and it is an unwise policy on the part of the producers of cream to insist on such low price for manufacturing that the creamerymen cannot afford to equip the creamery with modern appliances or to collect the cream at least three times a week. Cheap creamery equipment and cheap buttermakers may be very expensive in the end to cream producers. Both creamerymen and patrons should remember that a reputation for finest goods will ensure the highest current price and often a premium in addition. This enviable position can only be reached by every one doing his or her best and by having the closest co-operation and harmony in all the work relating to the creamery and the farm.

Summary of Important Notes.

For the Patron.

- 1. It pays to make cows comfortable at all times.
- 2. It pays to treat cows with invariable kindness. They should never be driven fast or worried by dogs.

app

- 3. Pure water should be provided for the cows, and they should be prohibited from drinking stagmant, impure water.
- 4. A box or trough containing salt, to which the cows have free access, should always be provided.
 - 5. Care must be taken to avoid feeds that will taint the milk.
- 6. The udders and flanks of the cows should always be washed or brushed clean before milking is commenced.
- 7. Milk from a freshly calved cow should not be skimmed until after the eighth milking.
 - 8. Only cream from cows in good health should be sent to the creamery.
 - 9. Tin pails only should be used.
- 10. Cream delivered every other day should be cooled as quickly as possible to 55 degrees and kept at that temperature or lower. If kept longer it should be cooled to under 50 degrees.
 - 11. Warm cream should never be mixed with cream already cooled.
 - 12. Every patron sending cream to a creamery should provide ice for cooling it
- 13. All vessels, including separator bowl, used in the handling of milk or cream should be thoroughly cleaned immediately after they are used by washing in lake

warm water and then thoroughly scalded with boiling water. A brush is preferable to a cloth for washing tinware or separators.

For Creamery Owners.

1. Pasteurizing the cream will give a better keeping quality of butter.

ream rately

hject

nable

it is

prices

with

mery

pro-

finest

This

id by

mer

riven

ibited

hould

clean

ighth

ble to

ng i: reatt luke

- 2. The use of a pure culture or starter in gathered cream will improve the keeping quality of the butter.
- 3. If the cream is not pasteurized, provide for cooling it quickly when delivered to the creamery.
 - 4. Provide an abundant supply of good, pure water for the creamery.
 - 5. Provide cold storage that will keep the but er at about 40 degrees or lower.
- 6. Support your buttermaker in dealing firmly with patrons who send cream which is not in good condition.

For the Buttermaker.

- 1. Attend personally, as far as possible, to the taking in, sampling and testing of the cream.
 - 2. Keep your creamcry clean, bright and tidy. Also yourself and assistants.
- 3. Be satisfied only with the finest quality of butter, the cleanest and the most attractive surroundings.

Copies of this bulletin may be obtained free for each patron of a creamery, by application to the Dairy and Cold Storage Commissioner, Ottawa.

